

DEPARTMENT OF MATHEMATICS & STATISTICS

MATH 2513

FINAL EXAMINATION  
DECEMBER 2000

TIME: 3 HOURS

**NO CALCULATORS PERMITTED**

MARKS

1. Given  $\vec{a} = (1, -1, 1)$ ,  $\vec{b} = (2, 3, 0)$  and  $\vec{c} = (5, 0, 12)$  calculate the following:

(3) (a) the cosine of the angle between  $\vec{b}$  and  $\vec{c}$ ;

(3) (b) the vector projection of  $\vec{a}$  onto  $\vec{c}$ ;

(3) (c)  $\vec{c} \cdot (\vec{a} \times \vec{b})$

(4) 2. (a) Find the parametric **and** symmetric equations of the line which is parallel to the line of intersection of the planes  $x + y = 2$  and  $-2x + z = 0$ , and which passes through the origin.

(3) (b) Find the equation of the plane which contains the point  $(1, 0, 0)$  and the line  $L : \frac{x-1}{2} = y+1 = \frac{z}{3}$ .

(4) 3. (a) Find the equation of the tangent plane to the surface  $x^2 + 4y^2 + 9z^2 = 25$  at the point  $(0, 2, -1)$ .

(4) (b) Find the directional derivative of

$$F(x, y, z) = \sin(xyz)$$

at the point  $(0, 1, -1)$  in the direction  $\vec{v} = (1, -1, 2)$ .

(4) (c) Use the chain rule to find  $\frac{\partial z}{\partial r}$  if  $z = x^2 - y^2$  with  $x = re^t$  and  $y = re^{-t}$ .

(5) 4. (a) Find and classify the critical points of

$$f(x, y) = x^3 - 6xy + 8y^3.$$

(5) (b) Use Lagrange multipliers to find the maximum and minimum value of  $f(x, y, z) = \frac{1}{x} + \frac{1}{y}$  subject to the constraint  $\frac{1}{x^2} + \frac{1}{y^2} = 1$ .

(5) 5. (a) Sketch the region  $R$  of integration bounded by  $y^2 = x^3$  and  $y = x$ , and compute  $\int \int_R xy \, dA$ .

(5) (b) Use polar coordinates to evaluate  $\int \int_R x \, dA$  where  $R$  is the region in the 2nd quadrant which is inside the circle  $x^2 + y^2 = 6y$ .

(4) 6. Use cylindrical coordinates to find the volume of the solid bounded by the cylinder  $x^2 + y^2 = 4$  and the planes  $z = 0$  and  $y + z = 3$ .

(4) 7. (a) Evaluate  $\int_C x \sin y \, dx + xyz \, dz$  along the arc  $C$  given by  
 $\vec{r}(t) = t\vec{i} + t^2\vec{j} + t^3\vec{k}$ ,  $0 \leq t \leq 1$ .

(4) (b) Show that

$$\vec{F}(x, y) = (2x + y^2 + 3x^2y)\vec{i} + (2xy + x^3 + 3y^2)\vec{j}$$

is conservative and find a function  $\phi$  such that  $\vec{F} = \vec{\nabla}\phi$ . Use  $\phi$  to evaluate  $\int_C \vec{F} \cdot d\vec{r}$  where  $C$  is the arc of an ellipse going from  $(0, 0)$  to  $(2, 1)$ .

---

(60)